

Poster-1-10

Pseudogap Suppression by Competition with Superconductivity in La-Based Cuprates

Julia Küspert,¹ Ron Cohn Wagner,¹ Chun Lin,¹ Karin von Arx,^{1,2} Qisi Wang,¹ Kevin Kramer,¹ Wojciech R. Pudelko,^{1,3} Nicholas C. Plumb,³ Christian E. Matt,^{1,3} Claudia G. Fatuzzo,⁴ Denys Sutter,¹ Yasmine Sassa,² Jiaqiang-Q. Yan,⁵ Jianshi-S. Zhou,⁶ John B. Goodenough,⁶ Sunseng Pyon,⁷ Tomohiro Takayama,⁷ Hidenori Takagi,⁷ Tohru Kurosawa,⁸ Naoki Momono,^{8,9} Migaku Oda,⁸ Moritz Hoesch,^{10,11} Cephise Cacho,¹⁰ Timur K. Kim,¹⁰ Masafumi Horio,¹ and Johan Chang¹

¹ Physik-Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

² Department of Physics, Chalmers University of Technology, SE-412 96 Göteborg, Sweden

³ Swiss Light Source, Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland

⁴ Institute of Physics, École Polytechnique Fédérale de Lausanne (EPFL), CH-1015 Lausanne, Switzerland

⁵ Materials Science and Technology Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, United States

⁶ Texas Materials Institute, University of Texas at Austin, Austin, Texas 78712, USA

⁷ Department of Advanced Materials, University of Tokyo, Kashiwa 277-8561, Japan

⁸ Department of Physics, Hokkaido University - Sapporo 060-0810, Japan

⁹ Department of Applied Sciences, Muroran Institute of Technology, Muroran 050-8585, Japan

¹⁰ Diamond Light Source, Harwell Campus, Didcot, OX11 0DE, United Kingdom

¹¹ DESY Photon Science, Notkestraße 85, 22607 Hamburg, Germany

We have carried out a comprehensive high resolution angle-resolved photoemission spectroscopy (ARPES) study of the pseudogap interplay with superconductivity in La-based cuprates. The three systems $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$, $\text{La}_{1.6-x}\text{Nd}_{0.4}\text{Sr}_x\text{CuO}_4$ and $\text{La}_{1.8-x}\text{Eu}_{0.2}\text{Sr}_x\text{CuO}_4$ display slightly different pseudogap critical points in the temperature versus doping phase diagram. We have studied the pseudogap evolution into the superconducting state for doping concentrations just below the critical point. In this setting, near optimal doping for superconductivity and in presence of the weakest possible pseudogap, we uncover how the pseudogap is partially suppressed inside the superconducting state. This conclusion is based on the direct observation of a reduced pseudogap energy scale and re-emergence of spectral weight suppressed by the pseudogap. All together these observations suggest that the pseudogap phenomenon is in competition with superconductivity for anti-nodal spectral weight.